

Randall S. Davis
Glynn Dennis, Jr
Mary R. Odom
Andrew W. Gibson
Robert P. Kimberly
Peter D. Burrows
Max D. Cooper

Fc receptor homologs: newest members of a remarkably diverse Fc receptor gene family

Authors' addresses

Randall S. Davis,
Division of Hematology/Oncology,
Peter D. Burrows, Max D. Cooper,
Division of Developmental and Clinical
Immunology,
Andrew W. Gibson, Robert P. Kimberly,
Division of Clinical Immunology and
Rheumatology,
Randall S. Davis, Andrew W. Gibson, Robert P. Kimberly,
Max D. Cooper,
Department of Medicine,
Mary R. Odom, Peter D. Burrows, Max D. Cooper
Department of Microbiology,
Max D. Cooper,
Departments of Pathology and Pediatrics, and
the Howard Hughes Medical Institute,
University of Alabama at Birmingham, AL,
Glynn Dennis, Jr,
Laboratory of Immunopathogenesis and
Bioinformatics, Clinical Services Program,
National Institute of Allergy and Infectious
Disease, Science Applications International
Corporation-Frederick, Frederick, MD, USA

Correspondence to:

Max D. Cooper
Howard Hughes Medical Institute
University of Alabama at Birmingham
WTI 378, 1824 6th Avenue South
Birmingham, AL 35294-3300, USA
Tel: +1 205 934 3379
Fax: +1 205 934 1875

Acknowledgments

The authors would like to thank Ms. Ann Brookshire for editorial assistance and Ms. Marsha Flurry for graphics expertise. This work has been supported in part by NIH grants AI39816 (M.D.C.), AI48098 (P.D.B.), and AR49084 (R.P.K.). RSD was supported by an NIH Hematology Training Grant, DK07488. M.D.C. is a Howard Hughes Medical Institute Investigator.

Immunological Reviews 2002

Vol 190: 123–136

Printed in Denmark. All rights reserved

Copyright © Blackwell Munksgaard 2002

Immunological Reviews
0105-2896

Summary: Newfound relatives of the classical Fc receptors (FcR) have been provisionally named the Fc receptor homologs (FcRH). The recent identification of eight human and six mouse FcRH genes substantially increases the size and functional potential of the FcR family. The extended family of FcR and FcRH genes spans ~15 Mb of the human chromosome 1q21–23 region, whereas in mice this family is split between chromosomes 1 and 3. The FcRH genes encode molecules with variable combinations of five subtypes of immunoglobulin (Ig) domains. The presence of a conserved sequence motif in one Ig domain subtype implies Ig Fc binding capability for many FcRH family members that are preferentially expressed by B lineage cells. In addition, most FcRH family members have consensus tyrosine-based activating and inhibitory motifs in their cytoplasmic domains, while the others lack features typical of transmembrane receptors. The FcRH family members, like the classical FcRs, come in multiple isoforms and allelic variations. The unique individual and polymorphic properties of the FcR/FcRH members indicate a remarkably diverse Fc receptor gene family with immunoregulatory function.

Introduction

Beginning with the initial identification of cellular immunoglobulin (Ig)-binding receptors more than 30 years ago, the field of Fc receptor (FcR) biology has evolved to a comprehensive understanding of the biological consequences resulting from the physical interactions of FcRs with the Fc portion of Ig (1–4). The early observations that linked antibodies with the Fc γ RI, Fc γ RII, Fc γ RIII, and Fc ϵ RI on effector cells led to the identification of complex regulatory networks that integrate innate immunity with the cell-mediated and humoral arms of adaptive immune responses (5–8). Characterization of the classical FcRs, encoded by genes in the human chromosome 1q21–23 region, includes the elucidation of genomic and amino acid sequences of multiple FcR isoforms, specific cellular expression patterns, signaling potential, polymorphic functional properties, and protein structures (9–14). Struc-

tural analysis has yielded insight into the physiochemical interactions required for Fc binding and the essential Ig binding domains that have been maintained throughout vertebrate evolution (15–20). Genetically engineered mice deficient in FcR genes have demonstrated the involvement of these receptors in autoimmune diseases and verified their key roles in hypersensitivity reactions (13, 21–28). While some facets of FcR function are still unresolved, it is abundantly clear that the classical FcRs play an important role in maintaining the intricate coordination and balance between cellular and humoral immunity in higher vertebrates.

Robust efforts to sequence the genomes of model organisms and the complementary analysis of expressed sequence tags (ESTs) derived from specialized tissues have led to the recent identification of many previously unrecognized genes. This information has enabled a detailed genetic analysis of the phylogenetic relatedness of large gene families. Comparative analysis of homologous genes (either orthologs or paralogs) in different organisms has shed light on their genomic relatedness and diversity and provided theoretical connections between syntenic regions and immune function. Complex loci conserved for ~500 million years of vertebrate evolution, including the major histocompatibility complex (MHC) and the rearranging B-cell and T-cell receptor gene families, are related not only by similar sequence and structure but also by shared functional relationships. Their co-evolution with other multigene families accounts for the remarkably integrated features of adaptive immunity. Such higher order relationships involving many different gene families are proving to be more common than anticipated. This is particularly true for members of the Ig-gene superfamily that encode the most common domain type in the human proteome (29, 30).

A large Ig-like gene family of *FcαR* relatives has been characterized in a human chromosome 19q13.4 region known as the leukocyte receptor cluster (LRC) of genes (31–36). This polymorphic gene family of ~26 members encodes transmembrane immunoreceptors with tyrosine-based activating and inhibitory signaling properties (37, 38). These include the *FcαR* (CD89) (39), Ig-like transcripts/leukocyte Ig-like receptors (ILT/LIR/MIR-CD85) (33, 40–42), natural killer Ig-like receptors (KIR/CD158) (43–46), leukocyte associated-inhibitory receptors (LAIR) (47), and NKp46 (48). The closest mouse relatives of the LRC encoded receptors are the paired Ig-like receptors (PIR) that reside in a mouse chromosome 7 region syntenic with the human chromosome 19q13.4 region (49). Interestingly, NKp46 is the only LRC member that has maintained orthologous properties in humans and mice (50). While the ILT/LIR genes are likely to

be the closest homologs of the murine PIRs (37, 51, 52), the KIRs, which have been identified in humans and other mammals but not in mice, are thought to represent a recent independent expansion of the primate LRC (38, 53, 54).

The ligands for the immunoreceptors encoded by LRC genes are typically other members of the Ig superfamily. Many LRC Ig-like receptors, including the KIRs and LIRs, have MHC class I and class I-like ligands (17, 41, 45, 55–61). Allelic variants of the KIR and LIR genes and haplotypic differences in the numbers of these genes are manifested by differences in the receptor repertoire between individuals and even between different subsets of cells within individuals (35, 62). While the functional consequences of the polymorphic extracellular regions of these receptors are not yet fully understood, their intracellular signaling capacity is conserved among family members; their possession of common activating or inhibitory cytoplasmic signaling motifs relates the LRC receptors to a larger group of paired Ig-like receptors that regulate the activation status of the cells that bear them (63, 64). The signaling properties that distinguish inhibitory isoforms reside in their cytoplasmic tails and bestow the ability to initiate cellular inhibition via immunoreceptor tyrosine-based inhibitory motifs (ITIM) (65–67). Activating isoforms may have long cytoplasmic tails with immunoreceptor tyrosine-based activation motifs (ITAM) or, more commonly, associate with ITAM-bearing adapter proteins via non-covalent interaction of charged amino acids in their respective transmembrane regions (68–73). The associated adapter proteins include DAP12 and DAP10/KAP10 (74–78), which are encoded by genes in the region surrounding the LRC on chromosome 19, and the Fc receptor common γ chain (*FcR γ C*) and CD3 ζ chains, the genes for which are located in the region surrounding the Fc receptor genes on chromosome 1 (79–84). Notable similarities between these two clusters of Ig-like receptor genes include genomic structure, amino acid content, Ig domain ligands, and tyrosine-based signaling, features that collectively suggest a shared phylogenetic background.

Identification of Fc receptor homologs

The large number of *FcαR* relatives in the LRC gene family suggested to us that an extended family of the classical Fc receptors might exist within the chromosome 1q21–23 region. Utilizing available sequence information, we identified an amino acid (aa) consensus sequence derived from the second domains of human *FcγRI* (CD64), *FcγRII* (CD32), *FcγRIII* (CD16), and the third domain of polymeric Ig recep-

tor. When this 32 aa motif was used in a Genbank database query, genomic clones from the 1q21–22 region were identified that contained five Ig superfamily members, which we have provisionally termed the Fc receptor homolog family (huFcRH1–5) (85, 86). Identification of ESTs in the Lymphochip database (87) inferred B-cell expression for many of these genes, an implication that is supported by RNA blot and reverse-transcriptase polymerase chain reaction (RT-PCR) analyses (85). The FcRH1–5 genes were independently identified as immunoglobulin superfamily receptor translocation associated genes (IRTA) through an analysis of the breakpoints of a t(1;14)(q21;q32) chromosomal translocation from a multiple myeloma cell line (88, 89). Others have identified members of this family as IFGP (IgSF, FcR, gp42) (90) and SPAP (SH2 domain-containing phosphatase anchor protein) genes (91). Further analysis with the FcR consensus motif and amino acid sequences, which include huFcRH1–5 specific Ig-like domains and the third Ig domain (D3) of huFcγRI, led to the identification of an Fc receptor related gene, huFcRX/FcRL/FREB, and its mouse ortholog, mFcRX (92–94). A similar database analysis has since identified additional novel FcRH family members in both humans and mice (see Fig. 1). The recognition of this extensive family of Fc receptor relatives reveals a previously unanticipated diversity for the Fc receptor family.

Genomic diversity of the Fc receptor cluster

In parallel with the LRC family members, the classical Fc receptors and the FcRHs are related by chromosomal proximity and genomic structure. The specific organization of these genes is provisional, but data from the public and private databases suggests the Fc Receptor Cluster (FRC) spans over 15 Mb in the human chromosome 1q21–23 region (Fig. 1). The FRC genes are flanked by the high affinity FcγRIα on the centromeric end and the ITAM bearing signaling chain CD3ζ on the telomeric end. The telomeric end of the cluster was initially characterized by linkage studies that mapped the location of the low affinity FcγRII/III genes to the 1q22–23 region, and studies identified their mouse orthologs in a syntenic region on mouse chromosome 1 (95–98). A more recent analysis of human genomic bacterial artificial chromosome (BAC) clones that overlap this region has clarified the specific gene locations within the low affinity locus (99). The newly defined human Fc receptor related genes, FcRX/FcRL/FREB and FcRY, are closely linked with the low affinity FcR genes and map within 40 kb of FcγRIIB. The other classical FcR genes, FcεRIα, and FcγRIα, and Fcγc, the gene for the FcR common gamma chain, reside centromeric of the

low affinity FcγRII/III genes. Between them lies another recently identified gene, FcRH6, which is positioned near a highly related pseudogene (ψ) that shares similar features (our unpublished data). The FcRH1–5 locus is approximately 2 Mb centromeric of FcεRIα, spans more than 300 kb, and is flanked at its telomeric end by a member of the cysteine rich scavenger receptor family, the 5pα gene. The specific position of FcγRI has recently been called into question because of differing placement by genome mapping projects. According to the latest approximation by the National Center for Biotechnology Information (NCBI) we have provisionally placed it in a location that correlates with its original positioning (100, 101).

Homology-based mapping of the human and mouse genomes identifies syntenic regions of human chromosome 1q21–23 that are split between mouse chromosomes 1 and 3. Although the exact boundary between these regions remains unclear, it appears to be centered near the CD1 locus. CD1A-E link within 800 kb centromeric of FcεRIα, which is located on mouse chromosome 1, and within 300 kb telomeric of 5pα, which resides on mouse chromosome 3 (102). Previous work indicates that CD1 defines a conserved linkage group border between human chromosome 1 and mouse chromosomes 1 and 3 (103, 104). The FRC region has been postulated to be among a group of paralogous MHC gene containing regions on chromosomes 1, 6, 9, and 19 that likely emerged through large-scale block duplications (105–107). The location of the genes for the CD1 MHC class I-like molecules, which play their main roles in innate immunity, in the midst of a family of genes that operate at the interface of innate and adaptive immunity, may portend related functions for these paralogous genes.

The 1q21–23 region is a hotspot for translocation events that have been defined in a number of human malignancies (108–111). Some of these translocations affect members of the FRC family, IRTA1/FcRH4 and FcγRIIB (89, 112). Given the location of the mouse chromosome 1 and 3 junction within the FRC and remarkable diversification of the extended Fc receptor family, it is tempting to draw a parallel between the breakpoint of paralogous MHC regions and involvement of these regions in the chromosomal instability and translocations recognized commonly in malignancy. Homology of intergenic elements in these paralogous regions, such as microsatellite repeats, retroelements, and other common repetitive sequences, potentially could provide a framework for promiscuous recombination, which can act as a driving force in natural evolutionary processes and malignant transformation (113–115).

The identification of novel members of the extended FcR

family on mouse chromosomes 1 and 3 helps to refine the definition of mouse and human FRC synteny. The portion of human chromosome 1q21–22 containing *FcγRIα* and *FcRH1–5* is located in a syntenic region of mouse chromosome 3 and is reversed with regard to its human genomic orientation. Characterization of the *mFcRHs* indicates considerable divergence relative to their human counterparts, particularly with respect to the number, order, and chromosomal organization of these genes as well as the location of the *Spα/CD5L/Ap16* gene within the *FcRH* locus. Despite inconsistent positioning of these genes, the close linkage of *Spα/CD5L/Ap16* in addition to the conserved genomic structure observed among *mFcRH1–3* (our unpublished data) indicates that these mouse genes reside in a syntenic location and share a common ancestry with their human FRC relatives. The location of *FcγRI* telomeric to *mFcRH1–3* not only suggests a common origin for this portion of the extended family, but, along with its linkage to the *FcRHs*, reaffirms the accuracy of its positioning on human chromosome 1q21–22. The location of *CD1D*, centromeric of the *mFcRH* genes, is consistent with its position

in humans near the junction between the two FRC derived regions.

The mouse chromosome 1 derived portion of the FRC locus is also in an opposite orientation relative to the syntenic region in humans, but it has generally conserved positioning of its homologous genes. The low affinity *FcγR* genes in this region have been recognized only as single copies in mice compared to their duplicate and diversified relatives in humans. The recent recognition of a third gene proximal to *mFcγRII* and *mFcγRIII*, *mFcrl3*, indicates another low affinity receptor may exist in mice, and we note that this receptor has greater identity to *huFcγRIII* (see ‘Phylogeny of the extended Fc receptor gene family’).

A particularly well conserved feature of FRC genes in both humans and mice is their possession of a split signal peptide encoded by two exons, the first of which most commonly contains both the 5'UTR and translation initiation start site (ATG). The second half of the signal peptide is consistently encoded by a 21 bp exon (97, 100, 116–119). This constitutes an important distinction between the FRC genes and




Fig. 1. Organization of the extended FRC gene family in humans and mice. The human 1q21–23 region is depicted along with its syntenic regions located on mouse chromosomes 1 and 3. Approximate positions were determined from NCBI, Celera, Ensembl, Mouse Genome Informatics (MGI) databases, and unpublished observations. Gene positions are approximations and are not to scale.

LRC genes, all of which have a 36bp second exon (120–123). The extracellular Ig-like domains of all FRC family members are encoded by single exons that follow the phase 1 splicing typical for Ig-like domains (9, 86). A notable exception is the FcRX gene, which does not maintain the 21 bp S2 exon, but rather its second exon encodes a partial Ig-like domain that also follows a phase 1 splicing pattern (92). Although the transmembrane and cytoplasmic regions differ widely among FRC family members, the exonic organization among FcRH genes is largely conserved (reviewed in 86). HuFcRH1-5 transmembrane regions are encoded by single exons that are followed by five exons encoding the cytoplasmic tails, the fifth of which includes the translation termination site and the 3'UTR. HuFcRH6 differs from huFcRH1-5 in that its cytoplasmic domain is encoded by four rather than five exons (our unpublished data). High-

lighting the diversity of the moFcRHs is the presence of a seventh exon in one moFcRH2 isoform that encodes a type B cysteine rich scavenger receptor domain. Notably, this is the first identification of an FRC gene encoding a chimeric molecule containing both Ig-like and cysteine rich scavenger receptor elements. The moFcRH2 scavenger domain is 56% identical to the amino terminal cysteine rich domain of the moSpα/CD5L/Ap16 scavenger receptor). This structural feature is not found in any of the huFcRHs or any currently characterized proteins in humans and mice. Other unique features of the extended family are found in the fifth exon of huFcRX and moFcRX, which encodes a proline and leucine rich domain not seen among other FcR or FcRH genes. This exon includes the translation termination site and the beginning of the 3'UTR. The classical FcR genomic structures have been described elsewhere (9, 10).




Fig. 2. Phylogenetic relationships of the extended Fc receptor gene family. The extracellular amino acids of each receptor were aligned using ClustalX, and the tree topology was estimated using neighbor-joining where branch values represent percentage bootstrap support after 500 replicates and values below 50% are not shown (47). The chromosome 19 encoded IgA-binding FcαR (CD89) was included in the analysis as a measure of tree topology.


Phylogeny of the extended Fc receptor family

The homology-based identification of multiple human FcR homologs indicates that the regulatory networks involved in modulating cellular function in response to FcR engagement may be surprisingly complex. Their shared chromosomal proximity, genomic structure, and Ig-like domains give credence to the conclusion that the FcR and FcRH gene families are modern descendants of an ancient genetic lineage. The conservation of key cytoplasmic elements involved in intracellular signaling cascades further infers complex biological functions, which may include the modulation of innate and adaptive immune responses. The recent identification of genes encoding additional glycoproteins, FcRH6, FcRX, and FcRY, that share sequence similarity and chromosomal proximity with the FcR and FcRH gene families indicates that this is a highly dynamic region of the mammalian genome. While Fc ϵ RI α , FcRX, and FcRY represent members of the FRC gene family that appear to maintain orthologous relationships in man and mouse, both the FcR and FcRH families appear to have undergone multiple rounds of gene duplication, exon duplication, and recombination since the speciation of rodent and primate lineages, thereby generating several paralogous receptors (118, 119, 124, 125).

Phylogenetic reconstruction of the extended Fc receptor family based on Ig-like domain amino acid sequences reliably segregates all known syntenic relatives into distinct FcR and FcRH families (Fig. 2). Included in this analysis is a novel mouse receptor with significant sequence similarity to huFc γ RIII. This receptor, termed moFcrl3 (NCBI Accession: NM_144559), shares a terminal node with huFc γ RIII that is segregated from the human and mouse Fc γ RII genes. Interestingly, the node containing huFc γ RIIA, huFc γ RIIB, and moFc γ RII also includes moFc γ RIII, thereby suggesting that the mouse gene currently designated as Fc γ RIII may actually represent an additional moFc γ RII gene. Human FcRH6, moFcRH6, ragp42, moFcRH1, and moFcRH2 segregate as

new members of the FcRH family, whereas the recently identified FcRX and FcRY relatives cluster with the classical FcRs. While this analysis cleanly separates the FcR and FcRH gene families into what may be considered distinct genetic lineages, the interdigitating chromosomal arrangement of these families indicates that the FcR and FcRH families may possess more complex relationships, possibly including genetic recombination between loci. Further evidence suggesting that the FcR and FcRH loci are actively involved in interlocus recombination comes from comparative analyses of individual Ig-like domains. We have shown that all Ig-like domains in the extended Fc receptor family can be grouped into five domain sub-types based on sequence similarity (85, 86) (Fig. 3A). Because it possesses all five sub-types of FcR/FcRH extracellular domain sequences, huFcRH3 was used in an all-against-huFcRH3 comparison of individual Ig-like domains to reaffirm the domain designations (Fig. 3B). Color-coding the extracellular domains of the extended Fc receptor gene family, based on sequence similarity to huFcRH3, depicts a high degree of similarity in domain architecture and highlights at least two possible examples of interlocus recombination events. Notably, domain 3 of huFcRH3 (yellow) represents a subunit that is present in all huFcRH family members, but is unique to huCD64 among the classical FcRs and contributes to its high affinity binding capacity of monomeric IgG (126–128) (see Fig. 3A). This type of Ig-like domain is also present in both FcRX and FcRY, which appear more closely related to the FcR family based on their chromosomal location and the phylogenetic reconstruction. The degree of genetic complexity emerging as a hallmark feature of the extended FcR gene family is reminiscent of the LRC-encoded multigene families (35), wherein homologous Ig-like receptors exhibit extensive extracellular sequence variation around a common Ig-like structure (59, 129). The core Ig-like structure of LRC encoded receptors is more closely related to the Ig-like domains of FcRs than any other known Ig domain structure and uniquely combines features of both the C2-set and I-set Ig domains (130). Identification of chicken Ig-like receptors (CHIR) that combine features of both the FRC and LRC suggests that the FRC and LRC genes may have belonged to a single gene family as recently as 250 million years ago. Avian homologs, like their mammalian counterparts, exist as paired receptors that share extracellular ligand binding potential but have opposing activating and inhibitory signaling potential. Penetration of these emergent properties into modern avian and mammalian representatives of this ancient genetic lineage attests to the importance of their biological function in vertebrate immunity. In keeping with this conjecture, chro-

Fig. 3. Conserved extracellular domain architecture. **A)** Ig-like domains identified in human and mouse members of the extended Fc receptor family are color-coded based on similarity to huFcRH3 domains. **B)** Heat-map depicting percentage similarities between individual huFcRH3 domains and all other Ig-like domains in the extended Fc receptor family. Sequence similarities were estimated from a Clustal generated all-against-huFcRH3 amino sequence alignment and domains were clustered based on their similarity measures (Spotfire Somerville, MA). Shaded cells represent the degree of similarity between the domain in that row and the FcRH3 domain above that column.



mosomal aberrations involving both the classical FcR and the FcRH gene families have been linked to autoimmunity and malignant transformation. Recently, an epistatic model of interaction between the LRC encoded KIRs and class I MHC alleles has been associated with delayed progression to AIDS in HIV-infected individuals (131).

The second domain of classical FcRs (color coded dark blue) is the major region of Ig–Fc interaction, and this domain clusters with homologous domains from 20 members of the extended Fc receptor family. Based on multiple sequence alignment of the domains designated as dark blue domains in Fig. 3A, we have identified an Fc-binding consensus sequence that distinguishes this domain from other domains in the extended Fc receptor family (Fig. 4, black shading). Notably, this consensus sequence does not include amino acid positions that have been shown to directly interact with Ig–Fc. Perhaps this finding reflects conservation of core residues that are essential for maintaining the structural requirements of Ig–Fc binding, while allowing for variation in the residues that confer ligand specificity (Fig. 4, yellow shading). The high degree of variability in the amino acid sequences of domains fitting the Ig–Fc binding consensus correlates with the decreased intensity displayed for the D2 cluster in the heatmap representation of domain similarity in Fig. 3B. Correspondingly, while D2 of moFcRL3 possesses the Ig–Fc binding consensus displayed in Fig. 4, the amino acid composition of this subunit is equally identical to Ig-like domain subtypes D2–D4, and thus it does not consistently cluster with other domains that match the Ig–Fc binding consensus (Fig. 3B). Interestingly, the identification of human cytomegalovirus encoded FcR homologs that bind Ig–Fc, despite their lack of the structural residues described in Fig. 4, indicates an unforeseen plasticity in the amino acid sequences that may bestow Ig-binding capability (132).

Diverse signaling potential of FcR family members

Members of the extended FcR family have diverse signaling capabilities based on their possession of tyrosine-based acti-

Fig. 4. Sequence similarity predicts Fc-binding potential for several members of the extended Fc receptor family. Sequences of Ig-like domains clustering with domain 2 (D2) of huFcγRIII and huFcεR were aligned using ClustalX. The original consensus sequence used to identify the human FcR homologs is shown atop the alignment (85). Residues shaded in black represent a consensus sequence that uniquely identifies this cluster of domains and yellow highlights the positions of residues that mediate Fc-binding for huFcγRIII (red letters) and huFcεR (green letters).

vation motifs and/or inhibition motifs that are shared by other activating and inhibitory pairs within the greater immunoreceptor family. The immunoreceptor tyrosine-based activation motifs (ITAM) contain two repeats of the consensus sequence Y-X-X-L/I spaced by 6–8 amino acids (E/D)-X-X-Y-X-X-(L/I)-X₆₋₈-Y-X-X-(L/I), while the immunoreceptor tyrosine-based inhibitory motifs (ITIM) feature a six amino acid consensus sequence (I/V/L/S)-X-Y-X-X-(L/V/I) (63–65, 67, 68). Following ligand binding by the activating receptor complexes, tyrosines in the ITAM are phosphorylated by src family kinases, enabling them to recruit other signaling elements in a signaling cascade that triggers cellular activation. In the case of ITIM bearing receptors, the phosphorylated tyrosines provide a docking site for phosphatases containing SH-2 domains that can abrogate cellular activation via signaling pathways that depend upon tyrosine phosphorylation (133, 134). The balance between activating and inhibitory receptor pairs can thus modulate cellular responses to a variety of stimuli.

The classical receptors FcγRI, FcγRIII (transmembrane form), and FcεRIα have short cytoplasmic tails and transduce activation signals via their non-covalent interaction with the single ITAM bearing FcRγc (79–82, 135, 136); additionally, FcγRIII can also transduce signals via the CD3ζ chain, which possesses three ITAMs (137, 138). The association of receptor subunits is dependent on a single aspartic acid residue present in the transmembrane regions of FcRγc, CD3ζ, FcγRIII, and FcεRIα. This charged residue is the key to the non-covalent interaction of these Fc binding molecules with ITAM bearing co-adaptor subunits. The association of FcγRI with the FcRγc subunit occurs via the former's basically charged histidine residue within the transmembrane region. Similar to FcγRIII and FcεRIα, the huFcRH1 transmembrane domain contains an acidic residue (glutamic acid) that could afford interaction with other transmembrane molecules. However, its possession of ITAM-like motifs in its cytoplasmic region suggests that huFcRH1 has autonomous signaling potential, and thus it differs from FcR relatives with short cytoplasmic tails that lack tyrosines. HuFcRH1 may thus be unique in the FcRH family in its possession of both types of immunoreceptor extracellular and adapter subunit characteristics.

In contrast with other classical Fc receptors, the receptors encoded by the FcγRII genes do not require co-association with adapter subunits to transduce intracellular signals. They instead possess either cytoplasmic ITAM sequences (FcγRIIA and FcγRIIC) or ITIM sequences (FcγRIIB) with corresponding activating or inhibitory functional capacity. The autonomous signaling property of FcγRII is shared by the FcRHs

that possess ITIMs, ITAMs, or both types of signaling motifs in their cytoplasmic tails (Table 1). The ITIMs found in the FcRH cytoplasmic tails maintain residues of the (I/V/L/S)-X-Y-X-X-(L/V/I) consensus motif. Tyrosine motifs present in some FcRH cytoplasmic domains share the previously defined ITAM consensus (E/D)-X-X-Y-X-X-(L/I)-X₆₋₈-Y-X-X-(L/I). For example, huFcRH1, huFcRH3, and moFcRH1 generally follow this defined motif. Other ITAM-like sequences are found among the FcRH cytoplasmic domains that maintain analogous positional conservation of tyrosine residues, but differ slightly in amino acid character at the +3 position. Although these sequences are highly reminiscent of ITAMs, functional studies will be required to clarify their functionality. Other tyrosine based motifs found in huFcRH1 and huFcRH3, may constitute 'hemi-ITAMs', wherein the sequence (E/D)-X-X-Y-X-X-(A/V) is present but a tandem second tyrosine-based sequence is not. A similar motif is noted in PIR-B (49).

Clearly the FcRHs are likely to differ substantially in their signaling capabilities relative to those of the FcR family members. Conversely, a signaling role is unlikely in the case of FcRX, since it does not possess a transmembrane region or N-linked glycosylation sites. Rather, its acidic biochemical characteristics are consistent with the likelihood that it functions intracellularly, perhaps as a scaffolding component interacting via its Ig-like domains. The evolutionary origins of the tyrosine-based functional capacities of the FcRHs versus those of its classical FcR relatives are not obvious. However, it is clear that similar tyrosine-based motif complexity is shared among FcRH family members. This may imply that the intracellular signaling elements engaged by these receptors will be shared among FcRHs, but may differ from those currently defined for other tyrosine-based immunoreceptors. The presence of both ITIM and ITAM in the same cytoplasmic tail adds complexity to their signaling potential. This signal transduction potential coupled with Ig-binding capacity of some FcRHs may involve new cellular signaling pathways.

Polymorphism of the Fc receptor family

The classical human receptors FcγRIIA and FcγRIIB are expressed in several distinct splice variant isoforms. In addition, FcγRI, FcγRII (A, B and C) and FcγRIII (A, transmembrane; and B, glycosylphosphatidylinositol (GPI)-anchored form) each have naturally occurring allelic variants that reflect non-synonymous single nucleotide polymorphisms (SNPs). The SNPs in the second extracellular domain (D2) of FcγRIIA and FcγRIIIA affect Fc binding affinity and specificity, while the

non-synonymous transmembrane SNP in Fc γ RIIB alters receptor signaling (R.P.K. unpublished results, 139). The short cytoplasmic tails of both Fc γ RIA and Fc γ RIIIA modulate signal transduction through non-covalent association with FcR γ c (140, 141), but the impact of the cytoplasmic sequence variants of Fc γ RIA and Fc γ RIIIA on the transduction of activation signals has not been established (142). Both the FcR γ c and CD3 ζ signaling partners of the FcRs appear to be invariant (143, 144).

Not surprisingly, members of the FcRH family may also display both splice variant and allelic diversity, the full extent of which has yet to be elucidated. Several splice variants have been reported that modify the open reading frame. These include two potentially secreted variants and a GPI anchored form of IRTA2 (FcRH5) (89) and a truncated form of FcRH2 (91). Multiple splice variants, including a D2-deficient sequence of FcRH4, have been identified on a message level (our unpublished results). Corresponding FcRH protein isoforms

have not yet been characterized, but the existence of at least some of these isoforms would parallel structural themes established for the classical FcRs. The frequency of SNPs in the FcRH coding regions is consistent with many other human genes (145). FcRH3 has one non-synonymous SNP in the D1 extracellular domain (Asn²⁸→Asp²⁸) and a second non-synonymous SNP in the cytoplasmic domain (Pro⁶⁶⁰→Leu⁶⁶⁰) that modifies a candidate ITAM motif. The latter SNP occurs with an allele frequency of 0.01 in Caucasians versus an allele frequency of 0.16 in the African-American population. This difference, coupled with similar allele frequencies in the two groups for the D1 SNP, is consistent with differential evolutionary pressure on the corresponding function. Unlike the classical FcR, however, FcRH3-5 do not appear to have SNPs that affect amino acid sequence in the second extracellular domain (A.W.G. and R.P.K. unpublished results). FcRH4 has a synonymous T>C transition at nucleotide 516 in D2 (Glu¹⁷²→Glu¹⁷²), but a second putative non-synonymous entry in

Table 1. Activating and inhibitory tyrosine-based signaling motifs

Receptor	ITAM-like	ITIM	Other tyrosines
huFcRH1	<u>E</u> FTY <u>L</u> NSPTPGQLQP <u>I</u> YENV <u>D</u> IY <u>S</u> RLRKANI TDVDY <u>E</u> DA		SGDEVY <u>S</u> LAY <u>N</u> QPEQ
huFcRH2	<u>E</u> FTY <u>S</u> SPTPDMEELQPVY <u>V</u> NV	<u>V</u> VY <u>S</u> QV <u>V</u> IY <u>S</u> S <u>V</u>	
huFcRH3	<u>E</u> PMY <u>S</u> NVNPGDSPNPIY <u>S</u> QI	<u>V</u> LY <u>S</u> EL	EDDEENYENVPRV
huFcRH4		<u>S</u> LYVDV <u>L</u> VY <u>S</u> EI <u>V</u> VY <u>S</u> EV	
huFcRH5	<u>E</u> PTYHNVPAWEELQPVYT <u>N</u> A	<u>V</u> VY <u>S</u> EV <u>L</u> IY <u>S</u> EV	
huFcRH6		<u>V</u> VY <u>S</u> VV	GEQCPLYANVHHQ
moFcRH1	<u>E</u> PLYENVNVVSGNEVY <u>S</u> LV		VLQG <u>S</u> TYPKSPDS QVSSGLY <u>S</u> KPRIN NIAHMDY <u>E</u> DAM
moFcRH3	<u>E</u> PTYYNVPACIELQPVY <u>S</u> NE	<u>V</u> IY <u>T</u> EV	
huCD32A	<u>D</u> GGY <u>M</u> T <u>L</u> NPRAPTDDDKNIY <u>L</u> T <u>L</u>		EETNNDY <u>E</u> TADGG
huCD32B		<u>I</u> TY <u>S</u> LL	ISALPGY <u>P</u> ECREM
moCD32B		<u>I</u> TY <u>S</u> LL	PEEVGEYRQPSGL(BI) PTSSSPYNPPDLE(BI) EETEHDY <u>Q</u> NHI
huFcR γ c	<u>D</u> GVYTGLSTRNQETY <u>E</u> T <u>L</u>		
moFcR γ c	<u>D</u> AVYTGLNTRSQETY <u>E</u> T <u>L</u>		
huCD3 ζ	<u>N</u> QLY <u>N</u> ELNLGRREEY <u>D</u> V <u>L</u> <u>E</u> GLY <u>N</u> ELQKDKMAEAY <u>S</u> E <u>I</u> <u>D</u> GLYQGLSTATKD <u>T</u> Y <u>D</u> AL		
moCD3 ζ	<u>N</u> QLY <u>N</u> ELNLGRREEY <u>D</u> V <u>L</u> <u>E</u> GVYNALQKDKMAEAY <u>S</u> E <u>I</u> <u>D</u> GLYQGLSTATKD <u>T</u> Y <u>D</u> AL		

Tyrosines and key positional residues of the ITAM and ITIM consensus motifs are underlined.

GenBank (C536T which changes Ser¹⁷⁹→Leu¹⁷⁹) has not been identified in the sequence analysis of FcRH4 in 48 African-Americans and 50 Caucasians.

The mouse ortholog of FcRH3 may also exist in polymorphic forms. Analysis of spleen mRNA from five inbred mouse strains indicates the existence of two alleles, one that is expressed in BALB/c, 129, CBA, and NZB mice and the other in C57BL/6 mice. The nucleotide discrepancies between the two forms result in peptide sequences that vary by the insertion or deletion of one residue and by single residue differences at nine other positions. These polymorphisms, located in the extracellular portion of the moFcRH3 protein, affect three of the four most amino-terminal Ig domains, namely D2, D3, and D4. One of the polymorphic residues creates a site for N-linked glycosylation in one allele only, and five of the residue differences, including the insertion, are situated among highly conserved residues in the portion of the peptide that is within the D2-Fc binding region of classical Fc receptors (dark blue domains Fig. 3A). The locations of these polymorphic residues suggest they could affect ligand binding.

Conclusion

The chromosome 1q21–23 region contains two related families of genes, the classical FcR (FcγRI, FcγRII, FcγRIII, and FcεRI) and the FcRH genes. The former have been shown to provide a structural basis for the molecular interaction between immune effector cells and the antibodies of the humoral immune system. This interaction results in phagocytosis, inflammatory responses, and the regulation of antibody production. The recent identification of the Fc receptor homologs promises to significantly expand this biological field of study. The diversity of structural features and signaling potentials of FcR/FcRH family members indicate an unanticipated complexity for the ligand-binding and functional repertoire of this extensive immunoreceptor family. The wealth of knowledge already accumulated for the classical FcRs will fuel the future biological definition of these new family members and broaden our understanding of the functional roles for this diverse family of molecules in host defense.

References

- Boyden SV, Sorkin E. The adsorption of antigen by spleen cells previously treated with antiserum *in vitro*. *Immunology* 1960;**3**:272.
- Berken A, Benacerraf B. Properties of antibodies cytophilic for macrophages. *J Exp Med* 1966;**123**:119–144.
- Henry C, Jerne NK. Competition of 19S and 7S antigen receptors in the regulation of the primary immune response. *J Exp Med* 1968;**128**:133–152.
- Paraskevas F, Lee ST, Orr KB, Israels LG. A receptor for Fc on mouse B-lymphocytes. *J Immunol* 1972;**108**:1319–1327.
- Ravetch JV, Clynes RA. Divergent roles for Fc receptors and complement *in vivo*. *Annu Rev Immunol* 1998;**16**:421–432.
- Heyman B. Antibody feedback suppression: towards a unifying concept? *Immunol Lett* 1999;**68**:41–45.
- Amigorena S, Bonnerot C. Fc receptors for IgG and antigen presentation on MHC class I and class II molecules. *Semin Immunol* 1999;**11**:385–390.
- Heyman B. Regulation of antibody responses via antibodies, complement, and Fc receptors. *Annu Rev Immunol* 2000;**18**:709–737.
- Ravetch JV, Kinet JP. Fc receptors. *Annu Rev Immunol* 1991;**9**:457–492.
- van De Winkel JG, Capel PJ. Human IgG Fc receptor heterogeneity: molecular aspects and clinical implications. *Immunol Today* 1993;**14**:215–221.
- Daeron M. Fc receptor biology. *Annu Rev Immunol* 1997;**15**:203–234.
- Kimberly RP, Salmon JE, Edberg JC. Receptors for immunoglobulin G. Molecular diversity and implications for disease. *Arthritis Rheum* 1995;**38**:306–314.
- Ravetch JV, Bolland S. IgG Fc receptors. *Annu Rev Immunol* 2001;**19**:275–290.
- Sondermann P, Kaiser J, Jacob U. Molecular basis for immune complex recognition: a comparison of Fc-receptor structures. *J Mol Biol* 2001;**309**:737–749.
- Sondermann P, Huber R, Oosthuizen V, Jacob U. The 3.2-A crystal structure of the human IgG1 Fc fragment-FcγRIII complex. *Nature* 2000;**406**:267–273.
- Garman SC, Würzburg BA, Tarchevskaya SS, Kinet JP, Jardetzky TS. Structure of the Fc fragment of human IgE bound to its high-affinity receptor FcεRIα. *Nature* 2000;**406**:259–266.
- Maxwell KF, et al. Crystal structure of the human leukocyte Fc receptor, FcγRIIa. *Nat Struct Biol* 1999;**6**:437–442.
- Garman SC, Kinet JP, Jardetzky TS. Crystal structure of the human high-affinity IgE receptor. *Cell* 1998;**95**:951–961.
- Hulett MD, Witort E, Brinkworth RI, McKenzie IF, Hogarth PM. Multiple regions of human FcγRII (CD32) contribute to the binding of IgG. *J Biol Chem* 1995;**270**:21188–21194.
- Tamm A, Schmidt RE. IgG binding sites on human Fcγ receptors. *Int Rev Immunol* 1997;**16**:57–85.
- Clynes R, Dumitru C, Ravetch JV. Uncoupling of immune complex formation and kidney damage in autoimmune glomerulonephritis. *Science* 1998;**279**:1052–1054.
- Nakamura A, et al. Fcγ receptor IIB-deficient mice develop Goodpasture's syndrome upon immunization with type IV collagen: a novel murine model for autoimmune glomerular basement membrane disease. *J F Exp Med* 2000;**191**:899–906.
- Yuasa T, et al. Deletion of Fcγ receptor IIB renders H-2(b) mice susceptible to collagen-induced arthritis. *J Exp Med* 1999;**189**:187–194.
- Kleinau S, Martinsson P, Heyman B. Induction and suppression of collagen-induced arthritis is dependent on distinct Fcγ receptors. *J Exp Med* 2000;**191**:1611–1616.

25. Bolland S, Ravetch JV. Spontaneous autoimmune disease in Fc(γ)RIIB-deficient mice results from strain-specific epistasis. *Immunity* 2000;**13**:277–285.
26. Dijkstra HM, van De Winkel JG, Kallenberg CG. Inflammation in autoimmunity: receptors for IgG revisited. *Trends Immunol* 2001;**22**:510–516.
27. Bolland S, Yim YS, Tus K, Wakeland EK, Ravetch JV. Genetic modifiers of systemic lupus erythematosus in Fc γ RIIB^{-/-} mice. *J Exp Med* 2002;**195**:1167–1174.
28. Takai T. Roles of Fc receptors in autoimmunity. *Nat Rev Immunol* 2002;**2**:580–592.
29. Lander ES, et al. Initial sequencing and analysis of the human genome. *Nature* 2001;**409**:860–921.
30. Venter JC, et al. The sequence of the human genome. *Science* 2001;**291**:1304–1351.
31. Kremer EJ, Kalatzis V, Baker E, Callen DF, Sutherland GR, Maliszewski CR. The gene for the human IgA Fc receptor maps to 19q13.4. *Hum Genet* 1992;**89**:107–108.
32. Dupont B, Selvakumar A, Steffens U. The killer cell inhibitory receptor genomic region on human chromosome 19q13.4. *Tissue Antigens* 1997;**49**:557–563.
33. Wagtmann N, Rojo S, Eichler E, Mohrenweiser H, Long EO. A new human gene complex encoding the killer cell inhibitory receptors and related monocyte/macrophage receptors. *Curr Biol* 1997;**7**:615–618.
34. Wende H, Colonna M, Ziegler A, Volz A. Organization of the leukocyte receptor cluster (LRC) on human chromosome 19q13.4. *Mamm Genome* 1999;**10**:154–160.
35. Wilson MJ, et al. Plasticity in the organization and sequences of human KIR/ILT gene families. *Proc Natl Acad Sci USA* 2000;**97**:4778–4783.
36. Trowsdale J, Barten R, Haude A, Stewart CA, Beck S, Wilson MJ. The genomic context of natural killer receptor extended gene families. *Immunol Rev* 2001;**181**:20–38.
37. Barten R, Torkar M, Haude A, Trowsdale J, Wilson MJ. Divergent and convergent evolution of NK-cell receptors. *Trends Immunol* 2001;**22**:52–57.
38. Volz A, Wende H, Laun K, Ziegler A. Genesis of the ILT/LIR/MIR clusters within the human leukocyte receptor complex. *Immunol Rev* 2001;**181**:39–51.
39. Maliszewski CR, March CJ, Schoenborn MA, Gimpel S, Shen L. Expression cloning of a human Fc receptor for IgA. *J Exp Med* 1990;**172**:1665–1672.
40. Cosman D, et al. A novel immunoglobulin superfamily receptor for cellular and viral MHC class I molecules. *Immunity* 1997;**7**:273–282.
41. Borges L, Hsu ML, Fanger N, Kubin M, Cosman D. A family of human lymphoid and myeloid Ig-like receptors, some of which bind to MHC class I molecules. *J Immunol* 1997;**159**:5192–5196.
42. Samaridis J, Colonna M. Cloning of novel immunoglobulin superfamily receptors expressed on human myeloid and lymphoid cells: structural evidence for new stimulatory and inhibitory pathways. *Eur J Immunol* 1997;**27**:660–665.
43. Lanier LL. NK cell receptors. *Annu Rev Immunol* 1998;**16**:359–393.
44. Brown MG, Scalzo AA, Matsumoto K, Yokoyama WM. The natural killer gene complex: a genetic basis for understanding natural killer cell function and innate immunity. *Immunol Rev* 1997;**155**:53–65.
45. Colonna M, Samaridis J. Cloning of immunoglobulin-superfamily members associated with HLA-C and HLA-B recognition by human natural killer cells. *Science* 1995;**268**:405–408.
46. Long EO, Wagtmann N. Natural killer cell receptors. *Curr Opin Immunol* 1997;**9**:344–350.
47. Meyaard L, et al. LAIR-1, a novel inhibitory receptor expressed on human mononuclear leukocytes. *Immunity* 1997;**7**:283–290.
48. Pessino A, et al. Molecular cloning of NKp46: a novel member of the immunoglobulin superfamily involved in triggering of natural cytotoxicity. *J Exp Med* 1998;**188**:953–960.
49. Kubagawa H, Burrows PD, Cooper MD. A novel pair of immunoglobulin-like receptors expressed by B cells and myeloid cells. *Proc Natl Acad Sci USA* 1997;**94**:5261–5266.
50. Biassoni R, Pessino A, Bottino C, Pende D, Moretta L, Moretta A. The murine homologue of the human NKp46, a triggering receptor involved in the induction of natural cytotoxicity. *Eur J Immunol* 1999;**29**:1014–1020.
51. Kubagawa H, et al. Paired immunoglobulin-like receptors of activating and inhibitory types. *Curr Top Microbiol Immunol* 1999;**244**:137–149.
52. Blery M, Kubagawa H, Chen CC, Vely F, Cooper MD, Vivier E. The paired Ig-like receptor PIR-B is an inhibitory receptor that recruits the protein-tyrosine phosphatase SHP-1. *Proc Natl Acad Sci USA* 1998;**95**:2446–2451.
53. Khakoo SI, et al. Rapid evolution of NK cell receptor systems demonstrated by comparison of chimpanzees and humans. *Immunity* 2000;**12**:687–698.
54. McQueen KL, Wilhelm BT, Harden KD, Mager DL. Evolution of NK receptors: a single Ly49 and multiple KIR genes in the cow. *Eur J Immunol* 2002;**32**:810–817.
55. Fanger NA, Cosman D, Peterson L, Braddy SC, Maliszewski CR, Borges L. The MHC class I binding proteins LIR-1 and LIR-2 inhibit Fc receptor-mediated signaling in monocytes. *Eur J Immunol* 1998;**28**:3423–3434.
56. Lanier LL. Follow the leader: NK cell receptors for classical and nonclassical MHC class I. *Cell* 1998;**92**:705–707.
57. Moretta A, et al. Receptors for HLA class-I molecules in human natural killer cells. *Annu Rev Immunol* 1996;**14**:619–648.
58. Chapman TL, Heikeman AP, Bjorkman PJ. The inhibitory receptor LIR-1 uses a common binding interaction to recognize class I MHC molecules and the viral homolog UL18. *Immunity* 1999;**11**:603–613.
59. Chapman TL, Heikema AP, West AP, Jr., Bjorkman PJ. Crystal structure and ligand binding properties of the D1D2 region of the inhibitory receptor LIR-1 (ILT2). *Immunity* 2000;**13**:727–736.
60. Boyington JC, Brooks AG, Sun PD. Structure of killer cell immunoglobulin-like receptors and their recognition of the class I MHC molecules. *Immunol Rev* 2001;**181**:66–78.
61. Fan QR, Garboczi DN, Winter CC, Wagtmann N, Long EO, Wiley DC. Direct binding of a soluble natural killer cell inhibitory receptor to a soluble human leukocyte antigen-Cw4 class I major histocompatibility complex molecule. *Proc Natl Acad Sci USA* 1996;**93**:7178–7183.
62. Shilling HG, et al. Allelic polymorphism synergizes with variable gene content to individualize human KIR genotype. *J Immunol* 2002;**168**:2307–2315.
63. Vely F, Vivier E. Conservation of structural features reveals the existence of a large family of inhibitory cell surface receptors and non-inhibitory/activatory counterparts. *J Immunol* 1997;**159**:2075–2077.
64. Ravetch JV, Lanier LL. Immune inhibitory receptors. *Science* 2000;**290**:84–89.
65. Daeron M, et al. The same tyrosine-based inhibition motif, in the intracytoplasmic domain of Fc γ RIIB, regulates negatively BCR-, TCR-, and FcR-dependent cell activation. *Immunity* 1995;**3**:635–646.
66. Daeron M, Vivier E. Biology of immunoreceptor tyrosine-based inhibition motif-bearing molecules. *Curr Top Microbiol Immunol* 1999;**244**:1–12.
67. Gergely J, Pecht I, Sarmay G. Immunoreceptor tyrosine-based inhibition motif-bearing receptors regulate the immunoreceptor tyrosine-based activation motif-induced activation of immune competent cells. *Immunol Lett* 1999;**68**:3–15.

68. Reth M. Antigen receptor tail clue. *Nature* 1989;**338**:383–384.
69. Bonnerot C, Amigorena S, Choquet D, Pavlovich R, Choukroun V, Fridman WH. Role of associated γ -chain in tyrosine kinase activation via murine Fc γ RIII. *EMBO J* 1992;**11**:2747–2757.
70. Cambier JC, Pleiman CM, Clark MR. Signal transduction by the B cell antigen receptor and its coreceptors. *Annu Rev Immunol* 1994;**12**:457–486.
71. Cambier JC. New nomenclature for the Reth motif (or ARH1/TAM/ARAM/YXXL). *Immunol Today* 1995;**16**:110.
72. Daeron M, Malbec O, Bonnerot C, Latour S, Segal DM, Fridman WH. Tyrosine-containing activation motif-dependent phagocytosis in mast cells. *J Immunol* 1994;**152**:783–792.
73. Weiss A, Littman DR. Signal transduction by lymphocyte antigen receptors. *Cell* 1994;**76**:263–274.
74. Lanier LL, Corliss BC, Wu J, Leong C, Phillips JH. Immunoreceptor DAP12 bearing a tyrosine-based activation motif is involved in activating NK cells. *Nature* 1998;**391**:703–707.
75. Campbell KS, Colonna M. DAP12: a key accessory protein for relaying signals by natural killer cell receptors. *Int J Biochem Cell Biol* 1999;**31**:631–636.
76. Olcese L, Cambiaggi A, Semenzato G, Bottino C, Moretta A, Vivier E. Human killer cell activatory receptors for MHC class I molecules are included in a multimeric complex expressed by natural killer cells. *J Immunol* 1997;**158**:5083–5086.
77. Wilson MJ, Lindquist JA, Trowsdale J. DAP12 and KAP10 (DAP10)-novel transmembrane adapter proteins of the CD3zeta family. *Immunol Res* 2000;**22**:21–42.
78. Chang C et al. Cutting edge: KAP10, a novel transmembrane adapter protein genetically linked to DAP12 but with unique signaling properties. *J Immunol* 1999;**163**:4651–4654.
79. Perez-Montfort R, Kinet JP, Metzger H. A previously unrecognized subunit of the receptor for immunoglobulin E. *Biochemistry* 1983;**22**:5722–5728.
80. Hibbs ML, et al. Mechanisms for regulating expression of membrane isoforms of Fc γ RIII (CD16). *Science* 1989;**246**:1608–1611.
81. Ernst LK, Duchemin AM, Anderson CL. Association of the high-affinity receptor for IgG (Fc γ RI) with the γ subunit of the IgE receptor. *Proc Natl Acad Sci USA* 1993;**90**:6023–6027.
82. Scholl PR, Geha RS. Physical association between the high-affinity IgG receptor (Fc γ RI) and the γ subunit of the high-affinity IgE receptor (Fc ϵ RI γ). *Proc Natl Acad Sci USA* 1993;**90**:8847–8850.
83. Pfefferkorn LC, Yeaman GR. Association of IgA-Fc receptors (Fc α R) with Fc ϵ RI γ 2 subunits in U937 cells. Aggregation induces the tyrosine phosphorylation of γ 2. *J Immunol* 1994;**153**:3228–3236.
84. Morton HC, et al. Functional association between the human myeloid immunoglobulin A Fc receptor (CD89) and FcR γ chain. Molecular basis for CD89/FcR γ chain association. *J Biol Chem* 1995;**270**:29781–29787.
85. Davis RS, Wang YH, Kubagawa H, Cooper MD. Identification of a family of Fc receptor homologs with preferential B cell expression. *Proc Natl Acad Sci USA* 2001;**98**:9772–9777.
86. Davis RS, Dennis G, Jr, Kubagawa H, Cooper MD. Fc receptor homologs (FcRH1-5) extend the Fc receptor family. *Curr Top Microbiol Immunol* 2002;**266**:85–112.
87. Alizadeh AA, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* 2000;**403**:503–511.
88. Miller I, Hatzivassiliou G, Cattoretti G, Mendelsohn C, Dalla-Favera R. IRTAs: a new family of immunoglobulin-like receptors differentially expressed in B cells. *Blood* 2002;**99**:2662–2669.
89. Hatzivassiliou G, et al. IRTA1 and IRTA2, novel immunoglobulin superfamily receptors expressed in B cells and involved in chromosome 1q21 abnormalities in B cell malignancy. *Immunity* 2001;**14**:277–289.
90. Guselnikov SV, et al. A family of highly diverse human and mouse genes structurally links leukocyte FcR, gp42 and PECAM-1. *Immunogenetics* 2002;**54**:87–95.
91. Xu MJ, Zhao R, Zhao ZJ. Molecular cloning and characterization of SPAP1, an inhibitory receptor. *Biochem Biophys Res Commun* 2001;**280**:768–775.
92. Davis RS, Li H, Chen CC, Wang YH, Cooper MD, Burrows PD. Definition of an Fc receptor-related gene (FcRX) expressed in human and mouse B cells. *Int Immunol* 2002;**14**:1075–1083.
93. Facchetti F, Cella M, Festa S, Fremont DH, Colonna M. An unusual Fc receptor-related protein expressed in human centroblasts. *Proc Natl Acad Sci USA* 2002;**99**:3776–3781.
94. Mechetina LV, et al. FCRL, a novel member of the leukocyte Fc receptor family possesses unique structural features. *Eur J Immunol* 2002;**32**:87–96.
95. Huppi K, Mock BA, Hilgers J, Kochan J, Kinet JP. Receptors for Fc ϵ and Fc γ are linked on mouse chromosome 1. *J Immunol* 1988;**141**:2807–2810.
96. Grundy HO, Peltz G, Moore KW, Golbus MS, Jackson LG, Lebo RV. The polymorphic Fc γ receptor II gene maps to human chromosome 1q. *Immunogenetics* 1989;**29**:331–339.
97. Qiu WQ, de Bruin D, Brownstein BH, Pearce R, Ravetch JV. Organization of the human and mouse low-affinity Fc γ R genes: duplication and recombination. *Science* 1990;**248**:732–735.
98. Oakey RJ, Howard TA, Hogarth PM, Tani K, Seldin MF. Chromosomal mapping of the high affinity Fc γ receptor gene. *Immunogenetics* 1992;**35**:279–282.
99. Su K, Wu J, Edberg JC, McKenzie SE, Kimberly RP. Genomic organization of classical human low-affinity Fc γ receptor genes. *Genes Immun* 2002;**3** (Suppl 1):S51–S56.
100. Osman N, Kozak CA, McKenzie IF, Hogarth PM. Structure and mapping of the gene encoding mouse high affinity Fc γ RI and chromosomal location of the human Fc γ RI gene. *J Immunol* 1992;**148**:1570–1575.
101. Dietzsch E, Osman N, McKenzie IF, Garson OM, Hogarth PM. The human FCG1 gene encoding the high-affinity Fc γ RI maps to chromosome 1q21. *Immunogenetics* 1993;**38**:307–309.
102. Shiina T, et al. Genomic anatomy of a premier major histocompatibility complex paralogous region on chromosome 1q21-q22. *Genome Res* 2001;**11**:789–802.
103. Moseley WS, Watson ML, Kingsmore SF, Seldin MF. CD1 defines conserved linkage group border between human chromosomes 1 and mouse chromosomes 1 and 3. *Immunogenetics* 1989;**30**:378–382.
104. Moseley WS, Seldin MF. Definition of mouse chromosome 1 and 3 gene linkage groups that are conserved on human chromosome 1: evidence that a conserved linkage group spans the centromere of human chromosome 1. *Genomics* 1989;**5**:899–905.
105. Lundin LG. Evolution of the vertebrate genome as reflected in paralogous chromosomal regions in man and the house mouse. *Genomics* 1993;**16**:1–19.
106. Flajnik MF, Kasahara M. Comparative genomics of the MHC: glimpses into the evolution of the adaptive immune system. *Immunity* 2001;**15**:351–362.
107. Kasahara M. The chromosomal duplication model of the major histocompatibility complex. *Immunol Rev* 1999;**167**:17–32.
108. Webber LM, Garson OM, Tate B, McKenzie IF, Hogarth PM. Fc receptor gene translocation in a t(1;19) pre-B ALL cell line. *Immunogenetics* 1990;**31**:356–360.
109. Williams DL, et al. New chromosomal translocations correlate with specific immunophenotypes of childhood acute lymphoblastic leukemia. *Cell* 1984;**36**:101–109.

110. Offit K, Wong G, Filippa DA, Tao Y, Chaganti RS. Cytogenetic analysis of 434 consecutively ascertained specimens of non-Hodgkin's lymphoma: clinical correlations. *Blood* 1991;**77**:1508–1515.
111. Cigudosa JC, et al. Cytogenetic analysis of 363 consecutively ascertained diffuse large B-cell lymphomas. *Genes Chromosomes Cancer* 1999;**25**:123–133.
112. Callanan MB, et al. The IgG Fc receptor, Fc γ RIIB, is a target for deregulation by chromosomal translocation in malignant lymphoma. *Proc Natl Acad Sci USA* 2000;**97**:309–314.
113. Dover G. Molecular drive: a cohesive mode of species evolution. *Nature* 1982;**99**:111–117.
114. Kulski JK, et al. The evolution of MHC diversity by segmental duplication and transposition of retroelements. *J Mol Evol* 1998;**46**:734.
115. Dawkins R, et al. Genomics of the major histocompatibility complex: haplotypes, duplication, retroviruses and disease. *Immunol Rev* 1999;**167**:275–304.
116. Kulczycki A, Jr, et al. Genomic organization of mouse Fc γ receptor genes. *Proc Natl Acad Sci USA* 1990;**87**:2856–2860.
117. van De Winkel JG, Ernst LK, Anderson CL, Chiu IM. Gene organization of the human high affinity receptor for IgG, Fc γ RI (CD64). Characterization and evidence for a second gene. *J Biol Chem* 1991;**266**:13449–13455.
118. Pang J, et al. Characterization of the gene for the human high affinity IgE receptor (Fc ϵ RI) α -chain. *J Immunol* 1993;**151**:6166–6174.
119. Ye ZS, Kinet JP, Paul WE. Structure of the gene for the α -chain of the mouse high affinity receptor for IgE (Fc ϵ RI). *J Immunol* 1992;**149**:897–900.
120. de Wit TP, Morton HC, Capel PJ, van De Winkel JG. Structure of the gene for the human myeloid IgA Fc receptor (CD89). *J Immunol* 1995;**155**:1203–1209.
121. Selvakumar A, Steffens U, Palanisamy N, Chaganti RS, Dupont B. Genomic organization and allelic polymorphism of the human killer cell inhibitory receptor gene KIR103. *Tissue Antigens* 1997;**49**:564–573.
122. Wilson MJ, Torkar M, Trowsdale J. Genomic organization of a human killer cell inhibitory receptor gene. *Tissue Antigens* 1997;**49**:574–579.
123. Alley TL, Cooper MD, Chen M, Kubagawa H. Genomic structure of PIR-B, the inhibitory member of the paired immunoglobulin-like receptor genes in mice. *Tissue Antigens* 1998;**51**:224–231.
124. Tepler I, et al. The gene for the human mast cell high-affinity IgE receptor α chain: chromosomal localization to 1q21-q23 and RFLP analysis. *Am J Hum Genet* 1989;**45**:761–765.
125. Le Coniat M, Kinet JP, Berger R. The human genes for the α and γ subunits of the mast cell receptor for immunoglobulin E are located on human chromosome band 1q23. *Immunogenetics* 1990;**32**:183–186.
126. Allen JM, Seed B. Isolation and expression of functional high-affinity Fc receptor complementary DNAs. *Science* 1989;**243**:378–381.
127. Hulett MD, Osman N, McKenzie IF, Hogarth PM. Chimeric Fc receptors identify functional domains of the murine high affinity receptor for IgG. *J Immunol* 1991;**147**:1863–1868.
128. Hulett MD, Hogarth PM. The second and third extracellular domains of Fc γ RI (CD64) confer the unique high affinity binding of IgG2a. *Mol Immunol* 1998;**35**:989–996.
129. Fan QR, Mosyak L, Winter CC, Wagtmann N, Long EO, Wiley DC. Structure of the inhibitory receptor for human natural killer cells resembles haematopoietic receptors. *Nature* 1997;**389**:96–100.
130. Dennis G, Jr, Kubagawa H, Cooper MD. Paired Ig-like receptor homologs in birds and mammals share a common ancestor with mammalian Fc receptors. *Proc Natl Acad Sci USA* 2000;**97**:13245–13250.
131. Carrington M, et al. HLA and HIV-1: heterozygote advantage and B*35-Cw*04 disadvantage. *Science* 1999;**283**:1748–1752.
132. Atalay R, et al. Identification and expression of human cytomegalovirus transcription units coding for two distinct Fc γ receptor homologs. *J Virol* 2002;**76**:8596–8608.
133. Unkeless JC, Jin J. Inhibitory receptors, ITIM sequences and phosphatases. *Curr Opin Immunol* 1997;**9**:338–343.
134. Long EO. Regulation of immune responses through inhibitory receptors. *Annu Rev Immunol* 1999;**17**:875–904.
135. Lin S, Cicala C, Scharenberg AM, Kinet JP. The Fc(ϵ)RI β subunit functions as an amplifier of Fc(ϵ)RI γ -mediated cell activation signals. *Cell* 1996;**85**:985–995.
136. Sutton BJ, Gould HJ. The human IgE network. *Nature* 1993;**366**:421–428.
137. Lanier LL, Yu G, Phillips JH. Co-association of CD3 ζ with a receptor (CD16) for IgG Fc on human natural killer cells. *Nature* 1989;**342**:803–805.
138. O'Shea JJ, Weissman AM, Kennedy IC, Ortaldo JR. Engagement of the natural killer cell IgG Fc receptor results in tyrosine phosphorylation of the ζ chain. *Proc Natl Acad Sci USA* 1991;**88**:350–354.
139. Stein MP, et al. C-reactive protein binding to Fc γ RIIa on human monocytes and neutrophils is allele-specific. *J Clin Invest* 2000;**105**:369–376.
140. Hou X, Dietrich J, Geisler NO. The cytoplasmic tail of Fc γ RIIIA α is involved in signaling by the low affinity receptor for immunoglobulin G. *J Biol Chem* 1996;**271**:22815–22822.
141. Edberg JC, et al. The cytoplasmic domain of human Fc γ RIa alters the functional properties of the Fc γ RI γ -chain receptor complex. *J Biol Chem* 1999;**274**:30328–30333.
142. Gibson AW, Wu J, Edberg JC, Kimberly RP. Fc γ receptor polymorphisms: Insights into pathogenesis. In: Kammer GM, Tsokos GC, eds. *Lupus: Molecular and cellular pathogenesis*. Totowa, NJ: Humana Press, 1999:557–573.
143. Wu J, Edberg JC, Gibson AW, Tsao B, Kimberly RP. Single-nucleotide polymorphisms of T cell receptor ζ chain in patients with systemic lupus erythematosus. *Arthritis Rheum* 1999;**42**:2601–2605.
144. Wu J, Edberg JC, Gibson AW, Kimberly RP. Conservation of Fc ϵ RI γ chain coding region in normals and in SLE patients. *Lupus* 2002;**11**:42–45.
145. Stephens JC, et al. Haplotype variation and linkage disequilibrium in 313 human genes. *Science* 2001;**293**:489–493.